

Lifestyle By Homecare Services

Pressure Ulcer Prevention Policy

**Health and Social Care Act
2008 (Regulated Activities) Regulations 2014** 12, 13, 17, 20

CQC Single Assessment Framework Topics

Safe Topic Areas:

Learning culture
Safe systems, pathways and transitions
Safeguarding
Involving people to manage risks
Safe environments
Safe and effective staffing
Infection prevention and control
Medicines optimisation

Effective Topic Areas:

Assessing needs
Delivering evidence-based care and treatment
How staff, teams and services work together
Monitoring and improving outcomes
Consent to care and treatment

Caring Topic Areas:

Kindness, compassion and dignity

Treating people as individuals
Responding to people's immediate needs

Responsive Topic Areas:

Person-centred care
Care provision, integration, and continuity
Listening to and involving people

Well-led Topic Areas:

Partnerships and communities
Learning, improvement and innovation

Please see the 'Quality Statements' section for full guidance

Scope

The scope of this policy is to provide guidance for all staff on:

- Strategies to prevent and reduce the risk of service users developing pressure ulcers,
- Escalation of concern to healthcare professionals,
- How to ensure best and evidence-based practice is implemented, and
- How to work in a multi-disciplinary team with healthcare professionals and other social care providers.

This policy and procedure are provided for the regulated activity of personal care.

Equality Statement

Our organisation is committed to equal rights and the promotion of choice, person centred care and independence. This policy demonstrates our commitment to creating a positive culture of respect for all individuals. The intention is, as required by the Equality Act 2010, to identify, remove or minimise discriminatory practice in the nine named protected characteristics of age, disability, sex, gender reassignment, pregnancy and maternity, race, sexual orientation, religion or belief, and marriage and civil partnership. It is also intended to reflect the Human Rights Act 1998 to promote positive practice and value the diversity of all individuals.

Key Points

- All staff within the organisation have a responsibility and role in prevention of pressure ulcers, to ensure pressure ulcer reduction strategies maintain a high profile.
 - All care staff should complete a body map for any mark, bruise, tear or pressure sore observed, this should be uploaded to the client's records/electronic records with accompanying notes and descriptions (see '[Appendix 2: Adult safeguarding decision guide, page 13](#)').
 - If new, staff should inform the Registered Manager to ensure a management plan is developed to prevent harm
 - An incident form will be completed for all identified category 2 - 4 pressure ulcers
 - Report multiple site grade 2, and individual site grade 3 – 4 pressure ulcers to CQC and safeguarding (use '[Appendix 1: Adult safeguarding decision guide, page 2](#)').
 - Grade 4 pressure ulcer require a full root cause analysis investigation.
 - The presence of a pressure ulcer creates a number of difficulties psychologically, physically and clinically to the service user, carers and family.
 - Pressure ulceration occurs when the skin and underlying tissues are compressed for a period of time, between the bone and the surface on which the service user is sitting or lying.
-
- A pressure ulcer occurs over a bony prominence.
 - It is an area of localised damage to the skin, and underlying tissue.
 - It can be caused by pressure, shear, friction and/or a combination of these.
-
- The GP will be informed that the service user has a pressure ulcer.
 - The organisation will fully comply with its duty under Regulation 20 the Duty of Candour, to act openly, honestly and to formally apologise where, through its, or its staffs' actions service users have come to harm, or could in the future, or have died.
 - There are a number of stages of pressure ulcers defined by the European Pressure Ulcer Advisory Panel Classification System (EPUAP).
-
- Stage I: non-blanchable
 - Stage II: partial thickness
 - Stage III: full thickness skin loss
 - Stage IV: full thickness tissue loss
 - Unstageable: full thickness skin or tissue loss
 - Suspected Deep Tissue Injury: Depth Unknown

Policy Statement

The following, from '[Pressure ulcers: revised definition and measurement, Summary and recommendations](#),' June 2018, provides indicative information about the nature of the problem: 'Pressure ulcers remain a concerning and mainly avoidable harm associated with healthcare delivery. In the NHS in England, 24,674 patients were reported to have developed a new pressure ulcer between April 2015 and March 2016, and treating pressure damage costs the NHS more than £3.8 million every day. Finding ways to improve the prevention of pressure damage is therefore a priority for policy-makers, managers and practitioners alike.'

Pressure ulcers are common in care settings and represent a significant burden of suffering for service users and carers, as well as being costly to treat. The presence of a pressure ulcer creates a number of difficulties psychologically, physically and clinically to the service user, carer and family. Pressure ulcer prevention and management should be person centred and an integral part of care assessment and planning, which requires a multi-disciplinary approach.

Please see '[Guidance Pressure ulcers: applying All Our Health.](#)' and see '[Safeguarding adults protocol: pressure ulcers and raising a safeguarding concern.](#)'

Pressure ulcers are preventable, and it is everyone's responsibility to reduce the risk of service users developing them whilst in their care.

The Policy

The organisation is committed to working in partnership with service users, carers, family and health care and social care professionals to minimise the risk of pressure ulcers and skin damage to service users.

This policy sets out the organisation's strategy for achieving this including:

- Responsibilities.
- Risk assessment and management.
- Strength-based assessment of need.
- Person-centred care planning.
- Identification and implementation of best and evidence-based practice.
- Staff training.
- Effective policies and procedures.
- Supporting nutrition and hydration.
- Escalation of needs to healthcare and social care professionals.
- Multi-disciplinary teamworking.

Responsibility

All staff within the organisation have a responsibility and role in prevention of pressure ulcers.

Please see '[Pressure ulcers, Skills for Care.](#)'

The Registered Manager

- Has overall responsibility for the safety of service users, ensuring we meet all the statutory
- Is to ensure pressure ulcer reduction strategies maintain a high profile.
- Is responsible for the development and implementation of training, policies and guidelines which are evidence-based and reflect best practice to support pressure ulcer prevention and
- Provides expert advice on pressure ulcer prevention and
- Ensures education and training is provided to all staff.
- Develops strategies to continuously reduce incidence of pressure
- Promotes a zero-tolerance approach to pressure ulcer development and ensure that any Grade 3 and 4 pressure ulcers are investigated using the Root Cause analysis (RCA) and action plans
- Uses '[Safeguarding adults protocol: pressure ulcers and raising a safeguarding concern to assess the need for referral to Safeguarding](#)' (March 2024) and '[Appendices 1 to 3: adult safeguarding decision guide, body map and concern proforma](#)' (March 2024).
- Reviews incidence data, observes trends and works with other local health professionals to reduce occurrence of pressure ulcers.
- Ensures pressure ulcer prevention equipment and resources are available and fit for purpose.
- Leads on the development of audit tools.

Staff

Staff must adhere to the policy:

- Only undertake activities which you are trained and competent to deliver, e.g. risk assessments.
- Ensure that you are trained in pressure ulcer prevention.
- Any skin damages/changes noted reported to the Manager or, if urgent, directly to the service user's GP.
- Any pressure ulcer or skin damage identified must be.

See '[Guidance Safeguarding adults protocol: pressure ulcers and raising a safeguarding concern.](#)'

Pressure Ulcer Risk Assessment and Management

This organisation employs PUP risk tool.

The primary assessment is the responsibility of the senior staff trained to undertake risk assessments, care assessment and care planning. All service users should be assessed using the pressure ulcer risk assessment tool.

Staff undertaking care assessments and reviews are trained in the use of the risk assessment tool, and the actions to follow where there are concerns identified.

Updated risk assessments will be undertaken when the service user's needs or condition changes that could increase their risk of a pressure ulcer including (but not exclusively):

- Significantly limited mobility (for example, people with a spinal cord injury),
- Significant loss of sensation,
- A previous or current pressure ulcer,
- Malnutrition,
- The inability to reposition themselves, or
- Significant cognitive impairment.

Additional factors that can significantly weaken skin and increase the risk of damage and developing a pressure ulcer include:


- Age 70+ as older people are more likely to have mobility problems and skin that's more easily damaged through dehydration and other factors.
- Moisture-associated skin damage (MASD)
 - Incontinence-associated dermatitis (IAD);
 - Intertriginous dermatitis (ITD), involving inflammation of the skin folds relating to perspiration;
 - Periwound MASD due to wound fluid leakage;
 - Peristomal MASD caused by leakage from stoma edges.
- Obesity.
- Medical conditions that affect blood supply, make skin more fragile or cause movement problems – such as:
 - Diabetes,
 - Peripheral arterial disease,
 - Kidney failure,
 - Heart failure,
 - Multiple sclerosis (MS) and
 - Parkinson's disease.

Further information can be found at ['Pressure Ulcers,' NHS](#).

Note: risk assessment tools should be used as an aide and not replace clinical judgement or negate staff concerns. For example, if a service user has gone from mobile to bedbound or chair bound, then risks will increase significantly and therefore extra vigilance on skin integrity and raising concerns to healthcare professionals is imperative, even if their risk assessment score still does not raise a concern.

Skin Assessment

- Skin inspection and assessment should occur regularly based on the risk score, instructions in the care plan, and or when needs change e.g. a service user's mobility significantly decreases or other risk factors develop.
- Assess skin condition of the service user immediately on admission to the service, especially bony prominences including sacrum, heels, hips, ankles, elbows, occiput and buttocks (diagram below).



- Record the results of the assessment and include within the care plan with relevant action that needs to be followed. Include areas of concern on a body map diagram.
- Any concern for skin integrity, areas of soreness or risks of pressure ulcer then the staff member must contact the office for escalation to the GP, or appropriate healthcare professional, or if urgent contact the GP directly.
- Where an existing skin and pressure ulcer management plan is in place this will be recorded within the assessment, including any treatment being provided e.g. creams and lotions. Prescribed medication for topological application will be recorded within the medication risk assessment and medication care plan.
- On initial assessment any areas of redness, concern or existing known or treated pressure sore damage will be recorded on a body map. All relevant information of the existing or treated skin damage or pressure ulcer will be recorded and included within the care and medication plan.
- Contact details of healthcare professionals treating the service user will be recorded within the assessment and care plan for advice and escalation as appropriate.
- Service users and carers who are willing or able, should be taught to assess their own (or family members) skin and take ongoing responsibility as appropriate to promote independence. The organisation will work in the best interests of the service user and in compliance with the Mental Capacity Act 2005.
- Service users and carers will be provided with a pressure ulcer prevention leaflet available from local NHS resources including Integrated Care Boards or Secondary Care.
- The condition of the skin will be assessed including:
 - Persistent erythema (reddening of the skin),

v1 Last Reviewed: Jodie Nolan Wed Apr 01 2026

Next Review: Registered Manager Sat Jan 16 2027

Locally Responsible:

- Non-blanching hyperaemia (redness which does not disappear on relief of pressure)
 - Blisters
 - Discolouration
 - Localised heat
 - Localised oedema (build-up of fluid in tissue and swelling)
 - Localised induration (loss of pliability and elasticity of skin)
- Identifying discolouration on service users with dark skin may be difficult and care should be taken not to rely solely on visual inspection. Erythema has traditionally been the main factor in recognising pressure damage, but staff need to consider the temperature, texture and presence of oedema to carry out a true skin assessment in service users with darkly pigmented skin.

Any concerns must be escalated to the service user's GP or other relevant healthcare professional through the office, or if urgent directly.

- Any existing or acquired pressure ulcers should be categorised using a validated assessment tool. See NICE '[Pressure ulcers](#),' last revised in March 2023. Suggested validation tools for adults include the [Braden Risk Assessment](#) tool, [Waterlow score](#), Norton risk assessment scale, or the [PURPOSE-T](#) (Pressure Ulcer Risk Primary or Secondary Evaluation Tool).

There are a number of stages of pressure ulcers defined by the panel:

- Stage I: non-blanchable.
- Stage II: partial thickness.
- Stage III: full thickness skin loss.
- Stage IV: full thickness tissue loss.
- Unstageable: full thickness skin or tissue loss.
- Suspected Deep Tissue Injury: Depth Unknown
- Where pressure damage is present, a comprehensive wound assessment will be completed. A tracing of the wound should be made and where possible, a photograph should be taken.
- Note: pressure ulcers should not be reverse/down categorised. For example, a grade 4 does not become a grade 2: this should be documented as 'Healing category 4.'

Please see '[Pressure ulcer categorisation, NHS](#).'

Categorisation of skin damage and pressure ulcers is a clinical decision made by an appropriately qualified healthcare practitioner and is provided for information to care staff.

Other Relevant Assessments

As part of the assessment process, new service users will also be risk assessed in the following areas, and outcomes from these will be used to support the pressure ulcer risk assessment process for that individual:

- Nutritional status and hydration
- Moving and handling assessment
- Pain assessment
- Continence assessment
- Mental health/capacity assessment

Documentation

Record details of the assessments in the service user's assessment documentation and care plan.

Ensure the date and time of the assessment is recorded and the information is signed by the assessing member of staff. Name and status should be written in block capitals.

Care provided will be evaluated and progress documented in the service user's care notes for each episode of care relating to skin care.

Where there is a pressure ulcer or skin integrity plan in place then information regarding application of medication, lotions and/or creams will be recorded in the care notes and/or the topical medication administration record (TMAR) or eMAR.

Where there is an identified pressure ulcer risk then the care plan will include actions to be taken at each visit, e.g. check skin integrity of heels, buttocks, elbows and apply cream/lotion in line with eMAR/TMAR.

Where instructions are provided from healthcare professionals on the management of skin conditions, this will be recorded within the care plan and eMAR/TMAR as required and actioned at each visit.

Where staff seek advice and input from healthcare professionals, this will be recorded in the care notes and plan, including times, date, health care professional name, role, telephone number and advice given. If advice is provided by email, this will be added to the care plan, and a note in the care notes instructing staff of the advice provided.

Care Plan

v1 Last Reviewed: Jodie Nolan Wed Apr 01 2026

Next Review: Registered Manager Sat Jan 16 2027

Locally Responsible:

Each service user will have in their care plan a pressure ulcer risk assessment, with any professional advice and actions required to ensure skin integrity and to minimise the risk of pressure ulcers.

All service users identified as at risk of pressure ulcer development (e.g. Waterlow score greater than 15), will have a written pressure ulcer prevention plan incorporating:

- Positioning and repositioning schedule
- Pressure relieving equipment
- Nutritional requirements
- Pain management
- Contenance management

This will be developed in partnership with the service user's GP, or other relevant healthcare professional managing their skin integrity, and incorporated within their care plan.

Where aids are provided to support the service user e.g. pressure care beds, staff will be provided with training to use these to effectively support the service user to minimise risks of pressure ulcers.

As part of the assessment process, and where additional aids may benefit the service user are identified, then signposting information will be provided to support the individual in seeking advice and accessing the aids.

Positioning

- Service users with an elevated risk and/or pressure ulcers, should be encouraged to actively mobilise, stand hourly (where it is safe to do so) and rest lying on their side for short periods in the day.
- Where possible, service users should be taught to reposition themselves and redistribute their weight and carers should be shown how to assist.
- Service users who are 'at risk' of pressure damage or have pressure damage, should be repositioned and the frequency of repositioning determined by the results of skin inspection and individual needs, not by a ritualistic schedule.
- Repositioning should take into consideration other relevant matters, including the service user's medical condition, their comfort, the overall plan of care, the support surface and attendance of formal and informal carers.
- Service users should be positioned in such a way as to minimise the impact on bony prominences and pressure ulcer.
- Service users should be positioned appropriately to reduce the effect of shear and friction forces.

A re-positioning schedule will be agreed with the service user and healthcare professionals and documented within the care plan. A repositioning chart or turning clock will be used as deemed necessary for individual service user's needs, e.g. poor continence, Waterlow greater than 21.

Seating

Service users are at a higher risk of pressure ulcer development when sitting out of bed if they are unable to mobilise and the seating is not designed for pressure management and or the aids being used are not matched to the service user's needs. Factors which increase risk are:

- Inability to reposition
- Incorrect/inappropriate chair or seating or aids
- Inability to redistribute weight
- Incontinence
- Poor nutrition and hydration

Where there are concerns for the service user identified within the risk management process (e.g. high risk identified within Braden Scale, Waterlow or Norton Risk Assessment), and no existing pressure ulcer management plan is in place, or the service user's needs have changed, then concerns will be escalated immediately to the service user's GP or other relevant healthcare professional via the office to request an assessment.

The staff assessing the service user will support in developing a pressure ulcer management plan. The referral to the GP or healthcare professional will include a request for review by an Occupational Therapist to identify aids to provide support, e.g. pressure management chair.

Heel Ulcer Prevention and Management

The incidence of pressure ulcer on the heels is high and need careful consideration and management as they can lead to prolonged medical intervention, infection and necrosis leading to amputation, and if septicaemia or gangrene occur even death.

Prevention of Heel Pressure

Service users should be encouraged to mobilise wearing good fitting footwear.

When service users are in bed or elevating the legs, the heels should be 'free-floating'. This can be achieved by the use of:

- Heel lifts, heel protectors, troughs and pillows placed lengthways.
- Anti-embolism stockings should be removed twice daily for a maximum of 30 minutes and skin inspected, in agreement with the GP practice or district nurse.
- The service user should be informed of when, how and frequency of removal of any compression, antiembolism stockings and importance of skin inspection by their GP practice or district nurse.

Moving and Handling

Skin damage can be minimised by using correct positioning, transferring and repositioning techniques and the use of aids.

For example: hoists, sliding sheets, pillows, bed cradles and other aids.

- Complete the service user moving and handling risk assessment within 24 hours of admission to the service.
- Write a plan of care to meet the service user's moving and handling needs.
- Service users should be encouraged to move independently where possible. If assistance is required, safer handling techniques should be employed. Refer to 'Moving & Handling Policy.'
- When hoisting service users, hoist slings must be the correct size and properly fitted. Hoist slings should not be left under service users.
- The use of four section electric profiling beds can contribute to reducing pressure, friction and shearing forces if the bed is used to its full potential:
 - Raise the end of the bed (reverse Trendelenburg).
 - Use the auto contour mechanism (raises the foot end, when head end is raised).
 - Use the knee-break facility to minimise sliding.
 - Remove slings, slide sheets or other parts of the handling equipment after moving the service user.
 - Do not use sheets to move service user, unless they are slide sheets.

Pain

- Service users who are in pain will be reluctant to reposition and relieve pressure.
- Service users' pain should be assessed individually, using local pain assessment tools and aids.
- A plan of care will be developed with the service user, so the pain is controlled and the service user is informed of the need for adequate analgesia.

- Staff will work with the service user and healthcare professionals to ensure effective pain relieve plan is in place.
- This will be detailed in the care plan and the Medication Administration Record.

Pressure Relieving Devices

These devices will be provided either privately by the service user or family, or by social services or NHS commissioners. Service users should be referred to social services for an assessment of needs and/or GP for referral to an Occupational Therapist.

Pressure relieving mattresses, cushions and devices do not replace the need for service user repositioning.

Mattresses

Pressure relieving support surfaces aim to reduce the magnitude and/or duration of pressure between the individual and support surface.

Choice of pressure relieving support surface should be made by a registered nurse or other healthcare professional who is trained and competent in pressure ulcer risk assessment, prevention and management.

Decisions on type of surface should be based on:

- Social assessment and support
- Pressure ulcer risk assessment
- Category (Grade) of pressure ulcer
- Ability to reposition
- Length of time spent out of bed
- Comfort
- Service user's weight and height
- General health
- Acceptability by the service user
- Requirement for bed side rails

For general guidance on equipment selection refer to the service equipment manual or the local product selection guides for specific information for clinical area.

Cushions

All service users who sit out of bed and are at a high risk of a pressure ulcer should be provided with a pressure relieving cushion or chair with integrated pressure relieving properties.

A specialist chair may require a referral to a physiotherapist or occupational therapist via primary care.

Safe Use of Equipment

When using equipment and aids to support the service user it is important to consider the impact of the equipment on other aspects of safety, e.g. falls.

Consider:

- Ensure mattress does not elevate the individual to an unsafe height where side rails are in place.
- Ensure the individual is within the recommended weight range for the equipment.
- All equipment should be cancelled and returned to the local authority or NHS provider as soon as it is finished with for maintenance and decontamination.
- **Remember:** when re-assessing service user's risk – think 'does the service user still require this equipment?'

Audit and Maintenance of Equipment

Equipment can deteriorate due to age and usage, therefore all pressure relieving equipment should be checked and maintained in good working order according to manufacturer's guidelines.

Refer to manufacturer's guidance on maintenance and decontamination processes in line with the 'Infection Prevention & Control Policy.'

Foam Mattress and Cushion Audit

Audits should be carried out regularly and all pieces of equipment should be checked for:

- Condition of the cover
- No stains
- Splits
- Tears

Electrical and mechanical equipment should be serviced in line with manufacturer warranty and guidance. Staff must not use equipment which is out of service period or which they believe to be

faulty or unsafe.

Where staff have concerns, they must immediately raise these to the office or manager who will follow up with the relevant organisation responsible for servicing the equipment, e.g. social services.

Maintain and Protect Skin Integrity

An individual's skin may be exposed to a variety of moist substances, which may make it more susceptible to shearing, friction and injury.

When moving and handling service users, all staff should take care not to damage a service user's skin and should:

- Not wear rings (other than wedding bands) or watches when turning or repositioning service users, and nails should be kept short and nail varnish removed.
- The skin should be kept well hydrated.
- If the service user is at high risk of skin damage or incontinent, they should be advised to use an emollient soap substitute (e.g. aqueous cream, emulsifying wax) to wash and apply moisturisers regularly.
- The service user's skin should be thoroughly dried using a patting motion, particularly over vulnerable areas. Do not use a rubbing motion or massage when drying as this causes friction forces and is associated with tissue damage.
- Talcum powder should not be used because of its tendency to cake, thereby increasing friction, clogging the pores and increasing risk of infection and skin damage.

Incontinence Management

All service users will be assessed for their continence needs, and a plan developed where required to support them.

- Incontinence should be managed effectively.
- Assess the service user and develop a plan of care.
- Request input from specialist incontinence service where appropriate.
- Barrier creams should be used with caution as they can clog the pores in the pad and effect efficacy.

Nutritional Status

v1 Last Reviewed: Jodie Nolan Wed Apr 01 2026

Next Review: Registered Manager Sat Jan 16 2027

Locally Responsible:

The following is extracted from 'Eat well, drink well and keep the skin well: Key nutrition and hydration messages to prevent pressure ulcers and promote wound healing, NHS Improvement January 2018.'

'Nutrition and hydration play a key role in keeping the skin healthy – so think EAT – evidence, assessment, take action – when assessing an individual's risk of developing a pressure ulcer (and the healing of existing pressure ulcers or wounds).'

Evidence

- Many nutritional risk factors have been identified in the development of pressure ulcers.
- Risk factors include low body mass index (BMI), malnutrition, nutritional deficiencies, unintentional weight loss, an impaired ability to eat independently and obesity.
- Obesity is an independent risk factor for developing a pressure ulcer, and the risk is further increased if the individual is obese and is malnourished.
- International guidance recommends using a nutritional screening tool to assess an individual's risk of malnutrition and also assessing their weight history, weight loss and ability to eat independently.
- The National Institute of Health and Care Excellence recognises nutrition deficiencies as a risk for developing pressure ulcers and supports the use of oral nutritional supplements for patients with identified nutritional deficiencies.
- Early identification and treatment of individuals who are malnourished or at risk of it are vital in preventing pressure ulcer development and promoting wound healing.

Assessment

- All service users should be screened on admission or first visit for nutrition and hydration risk, and reassessed regularly using assessment tools.
- All service users should receive a well-balanced diet in accordance with their wishes.
- All service users should have their Body Mass index (BMI) calculated on assessment or first visit where possible and repeated in line with risk assessed need.
- If the service user has a poor intake, a personalised food chart should be commenced.
- Service users with pressure ulcer/s should be referred to the dietician for further assessment through their GP if there are any concerns.
- Refer for specialist advice as required.
- Provide service users advice on well-balanced diet and protein-energy foods.
- Aim for two litres of fluid per day where this does not conflict with clinical advice and be respectful of service users' wishes.

Training and Education

All staff must read and follow this policy and will have training on pressure ulcer prevention, risk assessment and planning of care within their induction programme and annual update training.

A record of individual staff members' education and training will be maintained by the Registered Manager.

Staff will be assessed for competence at supervisions and spot checks to ensure they are competent in the management of pressure ulcer risk.

Care Staff

Once care staff have received training and are deemed competent, as recorded within their training file, they can:

- Assess risk of pressure ulcer
- Manage a Grade 1 pressure
- Document clearly and escalate any concerns and deterioration to the Manager, who will contact the appropriate clinician to seek guidance and support for the service user.

Safeguarding Adults and Skin Damage

Skin damage has a number of causes, some relating to the individual service users, such as poor medical condition, and others relating to external factors such as poor care or lack of resources (equipment or staffing).

It is recognised that not all skin damage can be prevented and therefore the risk factors in each case should be reviewed on an individual basis before a safeguarding referral is considered. Not all pressure ulcers in adults at risk are the result of neglect.

If there are concerns that observed pressure damage maybe the result of neglect or omissions care, please refer to the 'Safeguarding Adults Policy.'

Reporting of Pressure Ulcers

- All care staff are responsible for reporting pressure ulcers.

- All care staff should complete a body map for any ,mark, bruise, tear or pressure sore observed, this should be uploaded to the client's records/electronic records with accompanying notes and descriptions (see '[Appendix 2: Adult safeguarding decision guide, page 13](#)').
- If new, staff should inform the Registered Manager to ensure a management plan is developed to prevent harm.
- An incident form will be completed for all identified category 2 - 4 pressure ulcers.
- Report multiple site grade 2, and individual site grade 3 – 4 pressure ulcers to CQC and safeguarding ('[Appendix 1: Adult safeguarding decision guide, page 2](#)').
- An incident form will be completed for all identified category 2-4 pressure
- Report any development of pressure sores of grade 3 or above that develops after the organisation takes on the care of the service user. See guidance 18(2) '[Regulation 18: Notification of other incidents.](#)'
- A grade 4 pressure ulcer requires a full root cause analysis investigation.
- The service user's GP will be informed that the individual has a pressure ulcer.
- The organisation will fully comply with its duty under Regulation 20: Duty of Candour, to act openly, honestly and to formally apologise where through its, or its staff's, actions service users have come to harm, or could in the future, or have died.
- Safeguarding will be notified of the incident in line with local and national safeguarding guidance. For further guidance see '[Guidance Safeguarding adults protocol: pressure ulcers and raising a safeguarding concern.](#)'

Dissemination and Implementation

This policy will be introduced to each staff member on induction and made available through Access Policies & Procedures.

Process for Monitoring Compliance and Effectiveness

The following processes will be used to monitor the impact of the policy:

- Audit of safeguarding incidents
- Audits of accident and incidents
- Client complaints
- Client surveys
- Other feedback received

Themes and trends will be identified from the monitoring and compliance activities to identify learning and continuous improvement activities associated with the management of pressure ulcer

prevention.

This policy will be evaluated and reviewed annually to ensure that it meets current evidence and best practice, and is in line with current regulations and legislation.

See '[Pressure ulcers: revised definition and measurement, NHS Improvement.](#)'

COVID-19 Statement

Currently there are no COVID-19 rules or restrictions in the UK, though this could change if there is another major outbreak.

It is the responsibility of the Registered Manager and organisation to monitor the English Government guidelines for any changes to the guidance and law. See '[Adult social care: guidance.](#)'

In addition, there is guidance in place from NHS England, the UK Government and Health and Safety Executive on how care settings should minimise risk to both care workers and service users.

This advice should be incorporated into the risk management and health and safety policies, procedures, and systems of work where appropriate.

Please see:

- [COVID-19 restrictions replaced by public health advice, HSE](#)
- [Infection prevention and control in adult social care: acute respiratory infection](#)

Specific additional ACS Guidance (used in conjunction with 'Infection prevention and control in adult social care: acute respiratory infection'):

- [Infection prevention and control in adult social care settings](#)
- [Infection prevention and control: resource for adult social care](#)
- [Infection prevention and control: quick guide for care workers](#)

References and Further Reading

[Guidance Pressure ulcers: applying All Our Health](#)

[Guidance, Safeguarding adults protocol: pressure ulcers and raising a safeguarding concern](#)

[Pressure ulcers, NICE](#)

[Duty of Candour, CQC](#)

[Safeguarding Adults Protocol Pressure Ulcers and the interface with a Safeguarding Enquiry issued by Department of Health and Social Care](#)

[Pressure ulcers; prevention and management Clinical guideline CG179, NICE](#)

[Pressure ulcers, Quality standard \(QS89\), NICE](#)

[Nutritional support in adults \(QS24\), NICE](#)

[Waterlow](#)

[Braden Scale, Healthcare Improvement Scotland](#)

[PURPOSE-T, University of Leeds](#)

[Norton Pressure Sore Risk Assessment, Royal Commissions](#)

[Eat well, drink well and keep the skin well: Key nutrition and hydration messages to prevent pressure ulcers and promote wound healing, NHS Improvement January 2018](#)

[COVID-19 restrictions replaced by public health advice, HSE](#)

[Infection prevention and control in adult social care: acute respiratory infection](#)

[Infection prevention and control in adult social care settings](#)

[Infection prevention and control: resource for adult social care](#)

[Infection prevention and control: quick guide for care workers](#)

Quality Statements

Learning culture

We have a proactive and positive culture of safety based on openness and honesty, in which concerns about safety are listened to, safety events are investigated and reported thoroughly, and lessons are learned to continually identify and embed good practices.

Safe systems, pathways and transitions

We work with people and our partners to establish and maintain safe systems of care, in which safety is managed, monitored and assured. We ensure continuity of care, including when people move between different services.

Safeguarding

We work with people to understand what being safe means to them as well as with our partners on the best way to achieve this. We concentrate on improving people's lives while protecting their right to live in safety, free from bullying, harassment, abuse, discrimination, avoidable harm and neglect. We make sure we share concerns quickly and appropriately.

Involving people to manage risks

We work with people to understand and manage risks by thinking holistically so that care meets their needs in a way that is safe and supportive and enables them to do the things that matter to them.

Safe environments

We detect and control potential risks in the care environment. We make sure that the equipment, facilities and technology support the delivery of safe care.

Safe and effective staffing

We make sure there are enough qualified, skilled and experienced people, who receive effective support, supervision and development. They work together effectively to provide safe care that meets people's individual needs.

Medicines optimisation

We make sure that medicines and treatments are safe and meet people's needs, capacities and preferences by enabling them to be involved in planning, including when changes happen.

Assessing needs

We maximise the effectiveness of people's care and treatment by assessing and reviewing their health, care, well-being and communication needs with them.

Delivering evidence-based care and treatment

We plan and deliver people's care and treatment with them, including what is important and matters to them. We do this in line with legislation and current evidence-based good practice and standards.

How staff, teams and services work together

We work effectively across teams and services to support people. We make sure they only need to tell their story once by sharing their assessment of needs when they move between different services.

Monitoring and improving outcomes

We routinely monitor people's care and treatment to continuously improve it. We ensure that outcomes are positive and consistent, and that they meet both clinical expectations and the expectations of people themselves.

Consent to care and treatment

We tell people about their rights around consent and respect these when we deliver person-centred care and treatment.

Kindness, compassion and dignity

We always treat people with kindness, empathy and compassion and we respect their privacy and dignity. We treat colleagues from other organisations with kindness and respect.

Treating people as individuals

We treat people as individuals and make sure their care, support and treatment meets their needs and preferences. We take account of their strengths, abilities, aspirations, culture and unique backgrounds and protected characteristics.

Responding to people's immediate needs

We listen to and understand people's needs, views and wishes. We respond to these in that moment and will act to minimise any discomfort, concern or distress.

Person-centred care

We make sure people are at the centre of their care and treatment choices and we decide, in partnership with them, how to respond to any relevant changes in their needs.

Care provision, integration, and continuity

We understand the diverse health and care needs of people and our local communities, so care is joined-up, flexible and supports choice and continuity.

Listening to and involving people

We make it easy for people to share feedback and ideas or raise complaints about their care, treatment and support. We involve them in decisions about their care and tell them what's changed as a result.

Partnerships and communities

We understand our duty to collaborate and work in partnership, so our services work seamlessly for people. We share information and learning with partners and collaborate for improvement.

Learning, improvement and innovation

We focus on continuous learning, innovation and improvement across our organisation and the local system. We encourage creative ways of delivering equality of experience, outcome and quality of life for people. We actively contribute to safe, effective practice and research.

[Key questions and quality statements - Care Quality Commission](#)

Appendix: Pressure Care Monitoring Chart

KEY - Care delivered
 ✓ - Yes
 X - No (record why)

Name												
Frequency of care delivery (circle as appropriate) 1hrly 2hrly 3hrly 4hrly												
Date												
Time – record using 24-hour clock												
Surface												
Mattress appropriate (please state)												
Cushion appropriate (please state)												
Functionality/integrity check of equipment performed												
Skin Inspection												
All pressure areas checked												
Redness present Y/N												
Keep moving												
B	Right side											
E	Left side											
D	Back											
CHAIR												
Incontinence												
Urine												
Bowels												
Nutrition												
Diet (please state)												
Fluids (please state)												
Supplement(s) (please state)												
Initials												

v1 Last Reviewed: Jodie Nolan Wed Apr 01 2026
 Next Review: Registered Manager Sat Jan 16 2027
 Locally Responsible: